

Water quality analysis of River Panv Dhoi in reference to its physico-chemical parameters and heavy metals

D.R. Khanna, J. Ashraf¹, Beena Chauhan¹, R. Bhutiani, Gagan Matta and V. Singh

Received on : 10-05-2009

Accepted on : 15-05-2009

Abstract

This paper deals with the analysis of different water parameters of River Panv Dhoi which flows through Saharanpur district. It is a streamfed river and a tributary of Hindon. The sample collection was usually completed during morning hrs. between 8:00 AM to 10:00 AM. The parameters like Temperature, Turbidity, Conductivity, Total Solids (TS), Total Suspended Solids (TSS), Total Dissolved Solids (TDS), pH, Dissolved Oxygen (DO), Free Carbon dioxide (CO₂), Biochemical Oxygen Demand (BOD), Chemical Oxygen Demand (COD), Alkalinity, Total Hardness, Chloride (Cl⁻) and heavy metals like Lead, Zinc, Mercury and copper were analyzed.

Keywords:- Water, River, Panv Dhoi, Saharanpur, Heavy metals

Introduction

Water is the most precious commodity of life. It is not only the basic need for sustaining human life but also vital to all the segments of economic development. Water is being adversely affected, qualitatively and quantitatively by all kinds of human activities on land, in air and/or in water.

River Panv Dhoi flows through Saharanpur district. It is a streamfed river and a tributary of Hindon. This river originates near Shanklapuri Shiv Mandir of Panwarka, than it goes to Saharanpur. It is about 15 km in length and then it mixes with river Dhamola (Fig. 1). Metallic scraps, metallic dust, oil, paper and wooden products, besides domestic garbage appear to be the main items of the sewage. Main drain carries away some effluent from the factories and also from the residential colonies. This drain carries domestic sewage, which is poured into river Panv Dhoi. Therefore, it can be well considered that it carries a variety of pollutants of equally different in physico-chemical nature. A lot of work

has been done to evaluate the impact of human interferences on different water bodies by Khanna *et al.* (1997), Khanna *et al.* (2000), Khanna and Bhutiani (2003) and Khanna and Chugh (2004) and many more. It is very much essential to monitor the water quality of different water bodies present to asses their suitability for different purposes. Therefore the physico-chemical parameters of river Panv Dhoi were analysed for different seasons during 2002-2003.

Materials and Method

The samples for different parameters were analysed with the help of the procedure described by Welch (1948), APHA (1980), Mathur (1982), Ross (1983), Trivedi and Goel (1984) and Khanna (1993). The water samples were collected from five different sampling sites Shanklapuri Shiv Mandir (A), Makhraj Ka Pul (B), Laldas Ka Baada (C), Jogyan Pul (D) and Near Dhamola (E). The sample collection was usually completed during morning hrs. between 8:00 A.M. to 10:00 A.M. The parameters like Temperature, Turbidity, Conductivity, Total Solids (TS), Total Suspended Solids (TSS), Total Dissolved Solids (TDS), pH, Dissolved Oxygen

Author's Address

Department of Zoology, Gurukul Kangri University, Hardwar. ☒

¹Department of Zoology, M.S. College Saharanpur, (U.P.)




Fig. 1: Map of District Saharanpur showing the situation of Panv Dhoi River

(DO), Free Carbon dioxide (CO_2), Biochemical Oxygen Demand (BOD), Chemical Oxygen Demand (COD), Alkalinity, Total Hardness, Chloride (Cl) and heavy metals like Lead, Zinc, Mercury and copper were analyzed.

Results and Discussion

In an aquatic ecosystem the physico-chemical parameters are the main important factors responsible for the biotic healthiness of the system. They show their effect individually and also collectively. The values of different physico-chemical parameters in different seasons were tabulated in the Table 1-10 and are graphically represented in Fig. 2 to 11.

The temperature is one of the most important parameter in an aquatic environment. In the present study, a difference in the fluctuation of water temperature was observed 10.4 ± 1.5 (minimum) in the winter season and 28.0 ± 0.5 (maximum) in monsoon season. Annual average values of temperature varied between $15.6^\circ\text{C} \pm 4.5$ to $25.0^\circ\text{C} \pm 4.4$. Minimum annual average value of temperature was observed at sampling station A and maximum was observed at sampling Station E. The water temperature showed an upward trend from winter season to summer season followed by a downward from monsoon season onwards. A more or less similar trend has been observed in the river Yamuna by Chakrabarty *et al.* (1959) and in the Kallayi (John, 1976). Badola and Singh (1981) also reported similar trend in

the river Alaknanda. Same trend of temperature was observed by Khanna *et al.* (2001) in river Ganga at Haridwar.

The present study showed conductivity fluctuation of $65.10 \text{ mmhos/cm}^2 \pm 2.98$ in winter season to $336.78 \text{ mmhos/cm}^2 \pm 0.58$ in monsoon season. Annual average values of conductivity varied between $142.07 \text{ mmhos/cm}^2 \pm 104.39$ to $191.44 \text{ mmhos/cm}^2 \pm 127.32$. Minimum annual average value of conductivity was observed at sampling station E and maximum was observed at Sampling Station A. Identical results were observed by Raina *et al.* (1982) and Mittal and Sengar (1990) from various Indian rivers. Khanna *et al.* (2003) also observed similar trend of conductivity in Ganga river at Bulandshahar.

The water of the river Panv Dhoi becomes start turbid from summer season onward and in rainy season the water was highly turbid. The lowest turbidity was recorded ($19.78 \text{ JTU} \pm 1.00$) in winter season and highest ($1000.00 \text{ JTU} \pm 150.21$) noted in monsoon season. Annual average value of turbidity varied between $49.58 \text{ JTU} \pm 40.44$ to $908.46 \text{ JTU} \pm 101.05$ in which maximum average value observed at sampling station E and minimum at sampling station B. Similar pattern was also reported by Badola and Singh (1981), Dobriyal *et al.* (1983) in the hill streams of the Garhwal Himalaya, Khanna *et al.* (1997) in river Ganga, and Seth *et al.* (2000) in the river Ganga.

In the present investigation it was noted that the total solids were maximum in monsoon season ($2400.00 \text{ mg/l} \pm 226.63$), which were responsible for the turbidity in the river. The total solids were recorded minimum ($68.44 \text{ mg/l} \pm 1.36$) in winter season due to gradual sedimentation of the filterable residue. Annual average values of total solids varied between $179.53 \text{ mg/l} \pm 168.53$ to $1726.94 \text{ mg/l} \pm 651.22$. Minimum average value of total solids was observed at sampling station A and maximum at sampling station E. The highest value of total solids range between 1873 mg/l and 3573 mg/l reported by Kumar and Sharma (2002) in the river Krishni. Total solids cause ecological imbalance in the aquatic ecosystem by mechanical abrasive action. Similar trends were shown by Khanna and Chugh (2004) during study

of water quality of River Ganga at Haridwar. But Khanna and Singh (2000) found that total solids were maximum in summer in the water of Suswa River.

Minimum values of total dissolved solids were obtained during winter season ($48.33 \text{ mg/l} \pm 1.57$) and maximum values during monsoon season ($1100.00 \text{ mg/l} \pm 55.00$). Annual average value of total dissolved solids varied between $105.24 \text{ mg/l} \pm 13.10$ to $766.66 \text{ mg/l} \pm 351.18$, in which minimum average value was obtained from sampling station C and maximum from sampling station E. Same study was also done by Khanna and Chugh (2004) in the river Ganga at Haridwar.

Minimum values of total suspended solids were obtained during winter season ($20.11 \text{ mg/l} \pm 0.11$) and maximum values during monsoon season ($1300.00 \text{ mg/l} \pm 120.04$). Annual average value of total suspended solids varied between $57.85 \text{ mg/l} \pm 54.94$ to $960.26 \text{ mg/l} \pm 307.78$, in which minimum average value was obtained from sampling station A and maximum from sampling station E. Same study was also done by Khanna and Chugh (2004) in the river Ganga at Haridwar.

The Panv Dhoi river at Saharanpur showed high pH value (8.88 ± 0.19) in monsoon season which might be due to increase chemical load in the river. The minimum pH values (7.07 ± 0.26) were observed in winter season. The annual average values of pH varied between 7.32 ± 0.25 to 8.65 ± 0.37 . Highest annual average value was recorded at sampling station E and minimum at station A. It was recorded during this study that pH of the river Panv Dhoi is slightly alkaline in nature. Identical results were reported by Sangu and Sharma (1985) in the river Yamuna and Khanna *et al.* (1999) in Ganga river. Khanna *et al.* (2001) in the river Ganga at Haridwar and Khanna *et al.* (2003) in the river Ganga at Bulandshahar have also shown the alkaline nature of river.

The bio-chemical oxygen demand was observed maximum ($530 \text{ mg/l} \pm 15.98$) in monsoon season and minimum ($1.39 \text{ mg/l} \pm 0.17$) in winter season. The annual average value of biochemical oxygen demand ranged between $2.26 \text{ mg/l} \pm 0.82$ to $468.33 \text{ mg/l} \pm 80.98$. The minimum average value was found at sampling station

A and maximum at sampling station E. Highest annual average value of bio-chemical oxygen demand at sampling station E may be due to drainage of several small sewage drains into the river and runoff of sludgy, silted sewage during months of rainy season. Khanna & Singh (2000) noticed peak values during summer in Suswa river and Khanna *et al.* (1997) observed peak values in monsoon season in river Ganga.

Chemical Oxygen Demand (COD) represents chemically oxidizable load of organic matter in water. It was noted highest ($1240.00 \text{ mg/l} \pm 145.26$) in monsoon season and minimum ($4.69 \text{ mg/l} \pm 0.18$) in winter season. The annual average value of COD ranged between $5.62 \text{ mg/l} \pm 0.69$ to $1075.00 \text{ mg/l} \pm 220.17$ whereas least average value was found at sampling station B and maximum at sampling station E. Similar trends of COD have shown by Khanna *et al.* (2002, 2003) in the river Ganga and Khanna & Singh (2000) in Suswa River at Raiwala.

Maximum dissolved oxygen was recorded ($11.78 \text{ mg/l} \pm 0.25$) in the winter season. The minimum value of dissolved oxygen ($0.07 \text{ mg/l} \pm 0.01$) was observed in monsoon season. The annual average value of dissolved oxygen ranged between $1.68 \text{ mg/l} \pm 1.47$ to $9.61 \text{ mg/l} \pm 1.41$, whereas the minimum annual average value of DO was observed at sampling station E and maximum was observed at sampling station B. The dissolved oxygen reduced gradually from summer onward due to turbidity which retarded the photosynthetic activity of aquatic flora. The temperature showed an inverse relationship with the DO almost throughout the study. The cause of maximum dissolved oxygen in winter is due to reduced rate of decomposition by decreased microbial activity at low temperature (Strommer and Smock, 1989). Chopra *et al.* (1990), Gopal and Sah (1993) and Sharma (1999) also have got the same result and have opined that low temperature in winter increases the oxygen retaining capacity of water and solubility of O_2 in water. This trend was also observed by Badola and Singh (1981) in the river Alaknanda. Khanna (1993, 2001) and Khanna and Chugh (2004) has also reported the same trends in the river Ganga at Haridwar. Free carbon dioxide was observed maximum ($6.24 \text{ mg/l} \pm 0.96$) in monsoon season due to



higher turbidity and water temperature, but was recorded minimum ($0.07 \text{ mg/l} \pm 0.01$) in winter season. Annual average value of free carbon dioxide varied between $2.02 \text{ mg/l} \pm 1.95$ to $4.74 \text{ mg/l} \pm 1.56$ in which maximum average value obtained from sampling point E and minimum observed at sampling point A. Pahwa and Mehrotra (1966) and Ray *et al.* (1966) have reported that the Ganga river contains maximum free carbon dioxide in rainy season at Allahabad. Khanna *et al.* (1997) and Seth *et al.* (2000) have also reported the same trends in the river Ganga at Haridwar but Khanna and Singh (2000) observed maximum free carbon dioxide during summer in Suswa River at Raiwala, Dehradun.

Alkalinity of water is a measure of weak acid present in it and of the cations balanced against them (Sverdrup *et al.*, 1942). The highest concentration ($820.00 \text{ mg/l} \pm 55.27$) was observed in summer season and lowest ($35.00 \text{ mg/l} \pm 1.38$) in winter season. The annual average value of alkalinity varied between $50.02 \text{ mg/l} \pm 15.76$ to $781.66 \text{ mg/l} \pm 62.11$ in which maximum average value was obtained from sampling station E and minimum from sampling station A. Similar observation was also obtained by Khanna and Chugh (2004), Holden and Green (1960), Tallying and Rzoska (1967) and Abdin (1948).

Maximum values of total hardness were recorded in monsoon season ($390.82 \text{ mg/l} \pm 26.71$) and minimum was recorded in summer season ($83.78 \text{ mg/l} \pm 0.56$). The annual average values of total hardness ranged from $95.66 \text{ mg/l} \pm 11.71$ to $279.74 \text{ mg/l} \pm 17.27$ in which maximum average value is recorded from sampling station E and minimum at sampling station A. Khanna *et al.* (1993, 2003) and Mishra (2003) observed hardness in river Ganga at Hardwar and found more or less similar trends in their study.

The highest and lowest values of chloride were found in monsoon and winter season, $68.32 \text{ mg/l} \pm 4.56$ and $1.97 \text{ mg/l} \pm 1.66$ respectively. The annual average value of chloride varied between $5.51 \text{ mg/l} \pm 5.29$ to $59.67 \text{ mg/l} \pm 3.16$ in which maximum average value was recorded from the station E and minimum was at sampling station A. Chlorides are present in

sewage, sewage effluents and farm drainage. Significant levels of chloride were shown by many rivers like Yamuna (Sengar *et al.*, 1985); Tungbhadra (Reddy and Venkateswarlu, 1987); Jhelum (Raina *et al.*, 1984) and Kshipra (Mishra and Saxena, 1984). CPCB (2003) reported the value of chloride in between 14 to 51 mg/l during Ganga monitoring from Bithur, Kanpur to Sangam, Allahabad.

The minimum value $0.0638 \text{ mg/l} \pm 0.0072$ of lead was found from sampling station A and maximum $5.3975 \text{ mg/l} \pm 0.7123$ from sampling station E. The annual average of lead (Pb) ranged between $0.0702 \text{ mg/l} \pm 0.0076$ to $4.8452 \text{ mg/l} \pm 0.6109$. The minimum value $0.0214 \text{ mg/l} \pm 0.0022$ of copper was found from sampling station A and maximum $1.9774 \text{ mg/l} \pm 0.3996$ from sampling station E. The annual average value of copper varied between $0.0550 \text{ mg/l} \pm 0.0292$ to $1.5945 \text{ mg/l} \pm 0.3729$ in which minimum average value of copper was obtained at sampling station A and maximum average value at sampling station E.

Hg is not required even in small amount by any organisms. Virtually all metals, including the essential metal micronutrients, are toxic if exposure levels are sufficient high. The increased circulation of toxic metals in recent times resulted in the unavoidable build up of such toxic substances in the human food chain. The minimum value $3.1224 \text{ mg/l} \pm 0.0467$ of zinc was found from sampling station A and maximum $5.0012 \text{ mg/l} \pm 0.1814$ from sampling station E. The annual average value of Zinc ranged between $3.1651 \text{ mg/l} \pm 0.0528$ to $4.8695 \text{ mg/l} \pm 0.1763$ in which minimum average value of Zinc was obtained from sampling station A and maximum was recorded at sampling station E. The annual average value of Mercury varied between 0.0000 ± 0.0000 to $0.0005 \text{ mg/l} \pm 0.0006$. The minimum was found as nil at sampling station A, B, C and maximum ($0.0009 \text{ mg/l} \pm 0.0007$) at sampling station E in. The concentration of these metals at sampling station C, D and E give a highly misleading picture of the degree of metal pollution. Khanna *et al.* (2003) also reported heavy metals in water of Ganga river at Bulandshahar.



Table 1. Seasonal variation in physico-chemical parameters of River Panv Dhoi at sampling station A (2002-2003)

Physico-Chemical Parameters	2002-2003			
	Summer	Monsoon	Winter	Average
Temperature(°C)	18.0 ± 1.3	18.4 ± 1.3	10.4 ± 1.5	15.6 ± 4.5
Conductivity (µmhos/cm ²)	138.56 ± 0.40	336.78 ± 0.99	99.00 ± 0.56	191.44 ± 127.32
Turbidity (J.T.U.)	33.22 ± 0.36	120.28 ± 0.37	19.83 ± 0.45	57.73 ± 54.54
Total Solids (mg/L)	96.70 ± 1.35	373.45 ± 1.26	68.44 ± 1.36	179.53 ± 168.53
T.D.S. (mg/L)	64.13 ± 1.41	252.56 ± 1.48	48.33 ± 1.57	121.67 ± 113.62
T.S.S. (mg/L)	32.57 ± 0.04	120.69 ± 0.08	20.11 ± 0.11	57.85 ± 54.94
pH	7.30 ± 0.31	7.57 ± 0.29	7.07 ± 0.26	7.32 ± 0.25
BOD (mg/L)	2.30 ± 0.27	3.08 ± 0.35	1.39 ± 0.17	2.26 ± 0.82
COD (mg/L)	5.02 ± 0.29	7.36 ± 1.09	4.69 ± 0.18	5.69 ± 1.45
DO(mg/L)	9.52 ± 0.34	7.34 ± 1.11	11.78 ± 0.25	9.55 ± 2.18
Free CO ₂ (mg/L)	2.02 ± 0.29	3.98 ± 0.19	0.07 ± 0.01	2.02 ± 1.95
Alkalinity (mg/L)	66.67 ± 1.34	48.39 ± 1.22	35.00 ± 1.38	50.02 ± 15.76
Total Hardness (mg/L)	83.78 ± 0.56	107.20 ± 0.71	96.00 ± 0.85	95.66 ± 11.71
Chloride (mg/L)	2.96 ± 1.57	11.58 ± 1.93	1.97 ± 1.66	5.51 ± 5.29

Table 2. Seasonal variation in physico-chemical parameters of River Panv Dhoi at sampling station B (2002-2003)

Physico-Chemical Parameters	2002-2003			
	Summer	Monsoon	Winter	Average
Temperature(°C)	19.8 ± 1.6	20.2 ± 1.2	12.8 ± 0.7	17.6 ± 4.2
Conductivity (µmhos/cm ²)	138.15 ± 1.00	336.61 ± 31.12	99.58 ± 7.04	191.44 ± 127.18
Turbidity (J.T.U.)	33.34 ± 1.92	95.62 ± 0.63	19.78 ± 1.00	49.53 ± 40.44
Total Solids (mg/L)	97.75 ± 1.82	377.55 ± 14.51	70.47 ± 1.21	181.92 ± 169.95
T.D.S. (mg/L)	65.17 ± 1.74	254.57 ± 10.79	49.35 ± 4.98	123.03 ± 114.19
T.S.S. (mg/L)	32.58 ± 1.98	122.98 ± 2.03	21.12 ± 1.23	56.88 ± 55.79
pH	7.35 ± 0.90	7.59 ± 0.28	7.09 ± 0.05	7.34 ± 0.25
BOD (mg/L)	2.88 ± 0.82	3.18 ± 0.58	1.57 ± 0.35	2.54 ± 0.86
COD (mg/L)	5.52 ± 0.98	6.35 ± 1.67	4.98 ± 0.79	5.62 ± 0.69
DO(mg/L)	9.65 ± 0.72	8.18 ± 1.02	11.01 ± 0.58	9.61 ± 1.41
Free CO ₂ (mg/L)	2.08 ± 0.06	4.20 ± 0.12	1.24 ± 0.97	2.56 ± 1.52
Alkalinity (mg/L)	67.67 ± 1.72	49.08 ± 1.21	35.94 ± 0.78	50.89 ± 15.94
Total Hardness (mg/L)	84.01 ± 1.67	108.33 ± 2.61	96.94 ± 0.76	96.44 ± 12.19
Chlorides (mg/L)	3.15 ± 0.92	12.05 ± 0.93	2.37 ± 1.22	5.85 ± 5.37



Table 3. Seasonal variation in physico-chemical parameters of River Panv Dhoi at sampling station C (2002-2003)

Physico-Chemical Parameters	2002-2003			
	Summer	Monsoon	Winter	Average
Temperature(°C)	22.1 ± 0.7	24.2 ± 0.9	14.2 ± 0.6	20.2 ± 5.2
Conductivity (µmhos/cm ²)	111.26 ± 6.27	300.24 ± 15.25	85.20 ± 1.79	165.56 ± 117.35
Turbidity (J.T.U.)	250.00 ± 1.34	270.00 ± 9.50	210.00 ± 4.56	243.33 ± 30.55
Total Solids (mg/L)	176.22 ± 13.57	213.32 ± 9.23	170.26 ± 2.67	166.60 ± 23.33
T.D.S. (mg/L)	99.20 ± 5.42	120.28 ± 14.26	98.26 ± 6.21	105.24 ± 13.10
T.S.S. (mg/L)	77.02 ± 4.31	93.04 ± 2.74	74.00 ± 1.26	81.53 ± 10.23
pH	7.48 ± 0.18	7.62 ± 0.20	7.40 ± 0.10	7.50 ± 0.09
BOD (mg/L)	8.44 ± 0.63	11.16 ± 0.29	5.87 ± 0.12	8.42 ± 2.74
COD (mg/L)	26.30 ± 1.26	35.40 ± 1.04	22.65 ± 0.44	28.11 ± 6.56
DO(mg/L)	7.85 ± 0.27	6.56 ± 0.45	8.27 ± 0.89	7.56 ± 0.89
Free CO ₂ (mg/L)	2.21 ± 0.27	4.12 ± 0.39	1.75 ± 0.34	2.69 ± 1.25
Alkalinity (mg/L)	272.98 ± 9.02	261.55 ± 2.54	249.00 ± 1.04	261.17 ± 11.99
Total Hardness (mg/L)	221.00 ± 1.39	229.32 ± 4.58	215.51 ± 7.21	221.94 ± 6.96
Chloride (mg/L)	32.72 ± 1.85	39.39 ± 1.09	31.26 ± 0.77	34.45 ± 4.33

Table 4. Seasonal variation in physico-chemical parameters of River Panv Dhoi at sampling station D (2002-2003)

Physico-Chemical Parameters	2002-2003			
	Summer	Monsoon	Winter	Average
Temperature(°C)	26.0 ± 0.6	26.9 ± 0.8	17.0 ± 0.7	23.4 ± 5.5
Conductivity (µmhos/cm ²)	101.90 ± 1.02	274.78 ± 6.84	70.20 ± 2.22	148.96 ± 110.11
Turbidity (J.T.U.)	550.80 ± 9.95	650.70 ± 17.12	400.24 ± 6.27	533.93 ± 126.08
Total Solids (mg/L)	1536.42 ± 41.23	2315.26 ± 112.21	1030.50 ± 79.62	1627.40 ± 647.20
T.D.S. (mg/L)	755.62 ± 22.32	1015.28 ± 14.21	390.50 ± 7.96	720.46 ± 313.86
T.S.S. (mg/L)	780.80 ± 65.83	1300.00 ± 120.04	640.00 ± 29.32	906.93 ± 347.60
pH	8.60 ± 0.24	8.40 ± 0.15	8.00 ± 0.10	8.42 ± 0.40
BOD (mg/L)	315.00 ± 8.94	325.00 ± 10.06	205.00 ± 14.28	281.66 ± 66.58
COD (mg/L)	750.00 ± 45.21	950.00 ± 40.75	605.00 ± 20.29	768.33 ± 173.22
DO(mg/L)	4.90 ± 0.98	4.00 ± 0.34	5.10 ± 0.45	4.66 ± 0.56
Free CO ₂ (mg/L)	3.74 ± 1.22	5.36 ± 0.46	2.67 ± 0.94	3.93 ± 1.36
Alkalinity (mg/L)	715.00 ± 47.68	710.25 ± 22.98	708.00 ± 38.12	711.08 ± 3.57
Total Hardness (mg/L)	272.50 ± 6.74	294.00 ± 2.34	261.00 ± 41.08	275.83 ± 16.75
Chloride (mg/L)	58.37 ± 14.78	60.00 ± 8.98	53.12 ± 21.49	57.16 ± 3.59



Table 5. Seasonal variation in physico-chemical parameters of River Panv Dhoi at sampling station E (2002-2003)

Physico-Chemical Parameters	2002-2003			
	Summer	Monsoon	Winter	Average
Temperature (°C)	27.0 ± 0.7	28.0 ± 0.5	20.8 ± 0.6	25.0 ± 4.4
Conductivity (µmhos/cm ²)	100.20 ± 4.27	260.90 ± 8.42	85.10 ± 2.98	142.07 ± 104.39
Turbidity (J.T.U.)	925.22 ± 69.32	1000.00 ± 150.21	800.00 ± 22.34	908.46 ± 101.05
Total Solids (mg/L)	1590.82 ± 167.34	2400.00 ± 226.63	1100.00 ± 175.00	1728.94 ± 651.22
T.D.S. (mg/L)	800.00 ± 29.82	1100.00 ± 55.00	400.00 ± 12.05	766.66 ± 351.18
T.S.S. (mg/L)	880.82 ± 14.99	1300.00 ± 87.73	700.00 ± 63.29	960.26 ± 307.76
pH	8.85 ± 0.18	8.68 ± 0.19	8.22 ± 0.10	8.63 ± 0.37
BOD (mg/L)	500.00 ± 35.20	530.00 ± 15.98	375.00 ± 14.71	468.33 ± 80.98
COD (mg/L)	1180.00 ± 0.99	1240.00 ± 145.26	825.00 ± 89.74	1075.00 ± 220.17
DO (mg/L)	2.02 ± 0.24	0.07 ± 0.01	2.96 ± 0.19	1.68 ± 1.47
Free CO ₂ (mg/L)	4.88 ± 0.64	6.24 ± 0.96	3.12 ± 0.72	4.74 ± 1.56
Alkalinity (mg/L)	820.00 ± 55.27	815.00 ± 45.72	710.00 ± 41.39	781.66 ± 62.11
Total Hardness (mg/L)	276.40 ± 18.45	390.82 ± 26.71	252.00 ± 22.94	279.74 ± 17.27
Chloride (mg/L)	50.54 ± 2.50	68.32 ± 4.56	50.17 ± 6.04	59.67 ± 3.15

Table 6. Seasonal variation in Heavy Metals of river Panv Dhoi at sampling station A (2002-2003)

Heavy Metals	2002-2003			
	Summer	Monsoon	Winter	Average
Lead (Pb) (mg/L)	0.0880 ± 0.0072	0.0758 ± 0.0070	0.0636 ± 0.0072	0.0702 ± 0.0076
Copper (Cu) (mg/L)	0.0666 ± 0.0067	0.0740 ± 0.0069	0.0214 ± 0.0022	0.0559 ± 0.0292
Zinc (Zn) (mg/L)	3.1224 ± 0.0467	3.1488 ± 0.0613	3.2242 ± 0.0536	3.1851 ± 0.0528
Mercury (Hg) (mg/L)	0.0000 ± 0.0000	0.0000 ± 0.0000	0.0000 ± 0.0000	0.0000 ± 0.0000

Table 7. Seasonal variation in Heavy Metals of river Panv Dhoi at sampling station B (2002-2003)

Heavy Metals	2002-2003			
	Summer	Monsoon	Winter	Average
Lead (Pb) (mg/L)	0.0697 ± 0.0077	0.0760 ± 0.0072	0.0847 ± 0.0085	0.0711 ± 0.0073
Copper (Cu) (mg/L)	0.0739 ± 0.0284	0.0814 ± 0.0301	0.0324 ± 0.0324	0.0626 ± 0.0263
Zinc (Zn) (mg/L)	3.2258 ± 0.0295	3.2260 ± 0.0251	3.1810 ± 0.0231	3.2106 ± 0.0250
Mercury (Hg) (mg/L)	0.0000 ± 0.0000	0.0000 ± 0.0000	0.0000 ± 0.0000	0.0000 ± 0.0000



Table 8. Seasonal variation in Heavy Metals of river Panv Dhoi at sampling station C (2002-2003)

Heavy Metals	2002-2003			
	Summer	Monsoon	Winter	Average
Lead (Pb) (mg/L)	1.645 ± 0.3830	1.982 ± 0.4136	1.224 ± 0.2184	1.6170 ± 0.3797
Copper (Cu) (mg/L)	0.0872 ± 0.0181	0.0935 ± 0.0186	0.0649 ± 0.0161	0.0835 ± 0.017
Zinc (Zn) (mg/L)	3.6527 ± 0.2056	3.8928 ± 0.2132	3.4812 ± 0.2001	3.6755 ± 0.2067
Mercury (Hg) (mg/L)	0.0000 ± 0.0000	0.0000 ± 0.0000	0.0000 ± 0.0000	0.0000 ± 0.0000

Table 9. Seasonal variation in Heavy Metals of river Panv Dhoi at sampling station D (2002-2003)

Heavy Metals	2002-2003			
	Summer	Monsoon	Winter	Average
Lead (Pb) (mg/L)	3.2755 ± 0.0796	3.3972 ± 0.08120	3.2527 ± 0.0612	3.3084 ± 0.0776
Copper (Cu) (mg/L)	0.8475 ± 0.1761	0.9976 ± 0.1810	0.6474 ± 0.161	0.8308 ± 0.1756
Zinc (Zn) (mg/L)	4.2186 ± 0.01230	4.3270 ± 0.0112	4.1374 ± 0.0149	4.2443 ± 0.0732
Mercury (Hg) (mg/L)	0.0001 ± 0.00000	0.0002 ± 0.00000	0.0000 ± 0.0000	0.0001 ± 0.0000

Table 10. Seasonal variation in Heavy Metals of river Panv Dhoi at sampling station E (2002-2003)

Heavy Metals	2002-2003			
	Summer	Monsoon	Winter	Average
Lead (Pb) (mg/L)	4.9492 ± 0.6260	5.3975 ± 0.7123	4.1889 ± 0.5600	4.8452 ± 0.6109
Copper (Cu) (mg/L)	1.5737 ± 0.3631	1.9774 ± 0.3996	1.2324 ± 0.2998	1.5945 ± 0.3729
Zinc (Zn) (mg/L)	4.9384 ± 0.1796	5.0012 ± 0.1814	4.6692 ± 0.1616	4.8695 ± 0.1763
Mercury (Hg) (mg/L)	0.0002 ± 0.0004	0.0009 ± 0.0007	0.0004 ± 0.0005	0.0005 ± 0.0006

References

- Abdin, G., 1948. Physical and Chemical investigations relating to algal growth in the river Nile. *Cairo. Bull. Inst. Egypt*, 29: 20-24.
- Abiotic factors on the Phytoplanktonic population of A Pond. *Env. Cons. J.*, 1(1): 41-46.
- APHA., 1980. *Standard Methods for the examination of water and waste water*. American Public Health Association, 1015 fifteen street, NW Washington, 15: 1- 1134.
- Badola, S.P. and Singh, H.R., 1981. *Fish and fisheries of river Alaknanda. Proc. Nat. Acad. Sci.*, 15(B): 133-142.
- C.P.C.B., 2003. *A report on colour problem of river Ganga*, Central pollution Control Board, Zonal Kanpur. pp:1-9.
- Chakarbarty, R.D., Ray, P. and Singh, S.B., 1959. A quantitative survey of plankton and physiological



- conditions of the river Jamuna at Allahabad in 1954-1955. *Indian J. Fish.*, 6(1): 186-203.
- Chopra, A.K., Madhwal, B.P. and Singh, H.R., 1990. Abiotic variable and Prima productivity of river Yamuna at Naugaon, Uttar Kashi, Garhwal. *Indian J. Ecol.*, 17(1): 61-64.
- Dobriyal, A.K., Singh, H.R. and Bisht, K.L., 1983. Diurnal variation in hydrobiological parameter of two Hill streams of Garhwal Himalaya, India. Uttar Pradesh. *J. Zool.*, 3: 30-34.
- Gopal, B. and Sah, M., 1993. Conservation and management of rivers in India; Case study of the Yamuna. *Environmental Conservation*, 20(2): 243-253.
- Holden, J.M. and Green, J., 1960. Hydrobiology and plankton of the river Sokoto. *J. Anim. Ecol.*, 29(1): 65-84.
- John. V., 1976. Hydrobiological studies on the river Kallayi in Kerala. *Indian J. Fish.*, 23: 72-85.
- Khanna, D.R. and Bhutiani Rakesh., 2003. Ecological status of Sitapur Pond at Hardwar (U.A.) India. *Indian J. Environ & Ecoplan.*, 7(1): 175-178.
- Khanna, D.R. and Chugh, T., 2004. *Microbial Ecology: A study of river Ganga at Hardwar*, 1-273.
- Khanna, D.R. and Singh, R.K., 2000. Seasonal fluctuations in the plankton of Suswa River at Raiwala (Dehradun). *Env. Cons. J.*, (2&3): 89-92.
- Khanna, D.R., 1993. *Ecology and Pollution of Ganga River*. Ashish Publishing House, Delhi. pp: 1-241.
- Khanna, D.R., Badola, S.P. and Dobriyal, A.K., 1993. *Plankton Ecology of the river Ganga at Chandighat, Haridwar*. Advance in Limnology Ed. By H.R. Singh. Narendra Publishing House, New Delhi. 171-174.
- Khanna, D.R., Badola, S.P. and Malik, D.S., 1997. Population of green algae in relation to Physico-Chemical Factors of the river Ganga at Lal Ji Wala, Hardwar. U.P. *J. Zool.*, 17(3): 237-240
- Khanna, D.R., Bhutiani, Rakesh and Trivedi, Manoj., 2002. Impact of Paper Mill Effluent on some water parameters of Hindon river, at Saharanpur. *Him. J. Env. Zool.*, 16(1): 125-128.
- Khanna, D.R., Gautam, A. and Sarkar, P., 2001. Water quality of Bathing Ghats of river Ganga at Hardwar. National Conference on status of Indian Environment, Pre-conference proceeding, ASEA. Rishikesh, p-38.
- Khanna, D.R., Malik, D.S., Seth, T.R. and Rupendra., 1999. Correlation between abiotic factors and planktonic population in river Ganga at Rishikesh (U.P.) : *Eco. Sys. & Env.*, Ed. by D.R. Khanna, A. Gautam and A. Goutam, Published by ASEA, Rishikesh. pp: 69-76.
- Khanna, D.R., Singh Shakun., Gautam, A. and Singh, J.P., 2003. Assessment of water quality of river Ganga in District-Bulandshahar (U.P.) *India. J. Nat. Conservation*, 15(1): 167-165.
- Khanna, D.R., Gautam, Ashutosh., Chugh, Tarun and Sarkar, Praveen., 2000. Impact of abiotic factors on the phytoplanktonic population of a pond. *Env. Con. J.*, 1(1): 41-46.
- Kumar, N. and Sharma, R.C., 2002. Water quality of river Krishna with reference to Physico-chemical parameters. *J. Natcon*, 14(2): 273-297.
- Mathur, R.P., 1982. *Water and Waste Water testing*. Nem Chand and Bros. Publishers, Roorkee. pp: 1-54.
- Mishra, N. and Saxena, A.B., 1984. The effect of sewage with special reference to aquatic insects in the river Kshipra (India). *Int. J. Environ. Studies*, 23: 191-208.
- Mishra, S. and Joshi, B.D., 2003. Assessment of water quality with few selected parameters of River Ganga at Hardwar. *Him. J. Env. Zool.*, 17(2): 113-122.
- Mittal, S. and Sengar, R.M.S. (1990): Phytoplanktonic diversity in relation to certain physico-chemical characteristics of river water. *Proc. Nat. Acad. Sci. India*, 60(B), III.
- Pahwa, D.V. and Mehrotra, S.N., 1966. Observation on fluctuations in abundance of plankton in relation to certain hydrological conditions of river Ganga. *Proc. Nat. Acad. Sci.*, 36 B(2): 157-189.
- Raina, R., B.A. Subla and D.P. Zutshi., 1982. Water quality and plankton of Jhelum River. *Int. J. Ecol. Environ. Sci.*, 8: 11-17.
- Raina, V., Saha, A.R. and Ahmed, S.R., 1984. Pollution studies on river Jhelum-I : An assessment of water quality. *Indian J. Env. Hlth.*, 26: 187-210.
- Ray, P., Singh, S.B. and Sehgal, K.L., 1966. A study of some aspects of the river Ganga and Jamuna at Allahabad (U.P.) in 1958-59. *Proc. Nat. Acad. Sci. India*, 36 B(3): 235-272.
- Reddy, P.M. and Venkateshwarlu, V., 1987. Assessment of water quality and pollution in river Tungbhadra near Kurnool (A.P.). *J. Environ. Biol.*, 8(2): 109-119.
- Ross, Fedrick C., 1983. *Introductory Microbiology* Charles E. Merrill Publishing company. A Bell & Howell Company, Columbus, Ohio. pp: 1-615.
- Sangu, R.P.S., and Sharma, K.D., 1985. Studies on Water Pollution on Yamuna River at Agra. *Indian J. Environ. Hlth.*, 27(3): 257-261.



Sengar, R.M.S., Sharma, K.D. and Pathak, P.D., 1985. Studies on distribution of algal flora in polluted and non-polluted regions in Yamuna river at Agra (U.P.). *J. Indian Bot. Soc.*, 64: 365-376.

Seth, T.R., Khanna, D.R., Gautam, A., Chugh, T. and Sarkar, P., 2000. Temporal trends of phytoplanktonic diversity in the river Ganga at Hardwar. *Him. J. Env. Zool.*, 14: 129-134.

Sharma, A., 1999. *Limnological studies of Ban Ganga and distributional pattern of stream bottom fauna*. Ph. D. Thesis, University of Jammu.

Strommer, J.L. and Smock, L.A., 1989. Vertical distribution and abundance of invertebrates within the sandy

substrate of low gradient head water stream. *Fresh Water Biology*, 22: 263-274.

Sverdrup, H.H., Johnson, M.W. and Fleming, R.H., 1942. *The oceans, their Physics, and Chemistry and General Biology*. Prentice Hall, Inc, New York.

Talling, J.F. and Rzoska, J., 1967. The development of plankton in relation to hydrobiological regime in Blue Nile. *J. Ecol.*, 55: 636-662.

Trivedi, R.K. and Goel, P.K., 1984. *Chemical and Biological Methods for water pollution studies*. Karad : Environmental Publications. pp:1-251.

Welch, P.S., 1948. *Limnological methods*. The Blakiston Co. Philadelphia. pp: 1-381.

Fig. 2. Seasonal variation in physico-chemical parameters of River Panv Dhoi at sampling station A (2002-2003)




Fig. 3. Seasonal variation in physico-chemical parameters of River Panv Dhoi at sampling station B (2002-2003)




Fig. 4. Seasonal variation in physico-chemical parameters of River Panv Dhoi at sampling station C (2002-2003)




Fig. 5. Seasonal variation in physico-chemical parameters of River Panv Dhoi at sampling station D (2002-2003)




Fig. 6. Seasonal variation in physico-chemical parameters of River Panv Dhoi at sampling station E (2002-2003)




Fig. 9. Seasonal variation in Heavy Metals of river Panv Dhoi at sampling station C (2002-2003)




Fig. 7. Seasonal variation in Heavy Metals of river Panv Dhoi at sampling station A (2002-2003)




Fig. 10. Seasonal variation in Heavy Metals of river Panv Dhoi at sampling station D (2002-2003)




Fig. 8. Seasonal variation in Heavy Metals of river Panv Dhoi at sampling station B (2002-2003)




Fig. 11. Seasonal variation in Heavy Metals of river Panv Dhoi at sampling station E (2002-2003)

